

FLEXIBLE AFFINE CONES AND FLEXIBLE COVERINGS

MATEUSZ MICHAŁEK, ALEXANDER PEREPECHKO, AND HENDRIK SÜSS

ABSTRACT. We provide a new criterion for flexibility of cones over varieties covered by flexible affine varieties. We apply this criterion to prove flexibility of affine cones over secant varieties of Segre–Veronese embeddings and over certain Fano threefolds. We further prove flexibility of total coordinate spaces of Cox rings of del Pezzo surfaces.

Let X be an affine variety over an algebraically closed field $\mathbf{k}$ of characteristic zero. We consider actions of the additive group $\mathbb{G}_a = (\mathbf{k}, +)$ on X . The subgroup of $\text{Aut}(X)$ generated by all $\mathbb{G}_a$ -actions is called the *special automorphism group* of X and will be denoted by $\text{SAut}(X)$. We are interested in transitivity of the $\text{SAut}(X)$ -actions on the smooth points of X . Recall that an action of a group G on a set M is called *m-transitive* if for every two m -tuples $(x_1, \dots, x_m)$ and $(x'_1, \dots, x'_m)$ of pairwise distinct elements of M there exists $g \in G$ such that $g.x_i = x'_i$ for $i = 1, \dots, m$.

We have the following remarkable result.

Theorem 0.1 ([AFK⁺13, Theorem 0.1]). *Let X be an irreducible affine variety of dimension ≥ 2 . Then the following conditions are equivalent.*

- (i) *The group $\text{SAut}(X)$ acts transitively on the regular locus X_{reg} .*
- (ii) *The group $\text{SAut}(X)$ acts m -transitively on X_{reg} for every $m > 0$.*
- (iii) *The tangent space of every $x \in X_{\text{reg}}$ is spanned by tangent vectors to orbits of $\mathbb{G}_a$ -actions.*

If these equivalent conditions are fulfilled the affine variety X is called *flexible*. As examples of flexible varieties, let us mention affine cones over del Pezzo surfaces of degree 4 and 5 [Per13], over flag varieties, and affine toric varieties without torus factors [AKZ12].

Our main theorem provides a new criterion for flexibility of affine cones, see Section 1 for the proof.

Theorem 1.4. *Let Y be a projective variety covered by flexible normal open affine subsets U_i such that $Y \setminus U_i = D_i$, where D_i are effective divisors equivalent to a very ample divisor H . Then the affine cone $X = \text{AffCone}_H Y$ is flexible.*

Recall that a Segre–Veronese variety is an embedding of $\mathbb{P}^{d_1} \times \dots \times \mathbb{P}^{d_n}$ by the very ample line bundle $\mathcal{O}(s_1) \boxtimes \dots \boxtimes \mathcal{O}(s_n)$. Further, given a projective variety $X \subset \mathbb{P}^n$, the Zariski closure of the union of the secant (resp. tangent) lines to X is called a *secant* (resp. *tangential*) variety of X . As the first application of Theorem 1.4, We deduce the following result, see Section 2 for details.

2010 *Mathematics Subject Classification.* 14R20, 14J50.

Key words and phrases. Automorphism group, transitivity, flexibility, affine cone, Cox ring, Segre–Veronese embedding, secant variety, del Pezzo surface.

The research of M. Michałek was supported by IP grant 0301/IP3/2015/73 of the Polish Ministry of Science. The research of A. Perepechko was carried out at the IITP RAS at the expense of the Russian Foundation for Sciences (project no. 14-50-00150).

Theorem 2.21. *Let $X = v_{s_1}(\mathbb{P}(V_1)) \times \cdots \times v_{s_n}(\mathbb{P}(V_n))$ be a Segre–Veronese variety. Then the affine cone over the secant variety of X is flexible. Further, if $s_1 = \cdots = s_n = 1$, then also the affine cone over the tangential variety of X is flexible.*

Our proof technique relies on linearization of the affine charts of the ambient projective space by triangular transformations. They are inspired by algebraic statistics, precisely by computation of cumulants [CCM⁺14, MOZ15, SZ13, Zwi12, Zwi15].

Section 3 contains preliminaries on smooth rational T -varieties of complexity 1. These are varieties X with an effective action of a torus T , where $\dim X = \dim T + 1$. Section 4 is devoted to flexibility of affine cones over such varieties.

In [APS14] it was shown that smooth varieties of this type admit a toric covering and for certain cones over these varieties we, indeed, obtain flexibility. For example, this applies to all known Fano threefolds with 2-torus action. We use below the list of Fano threefolds in Mori–Mukai’s classification [MM81].

Theorem 4.5. *All affine cones over the Fano threefolds Q , 2.29, 2.30, 2.31, 2.32, 3.8, 3.18, 3.19, 3.23, 3.24, 4.4, and elements of the families 2.24, 3.10 admitting a 2-torus action in Mori–Mukai’s classification are flexible.*

The main tool to obtain these results is the combinatorial description of T -varieties developed in [AH06, AHS08], which in the case of complexity 1 allows to study (torus equivariant) coverings as in Theorem 1.4.

While Theorems 2.21 and 4.5 are concerned with projective coordinate rings, in Section 5 of the paper we obtain related results for total coordinate rings or Cox rings.

Theorem 5.4. *The total coordinate spaces of smooth del Pezzo surfaces are flexible.*

This was known so far only for the toric del Pezzo surfaces (i.e. those of degree 9, 8, 7 and 6) and by [AKZ12, Thm. 0.2] for the case of degree 5, where the total coordinate space is known to be the affine cone over the Grassmannian $G(2, 5)$. On the other hand, this extends a result of [APS14], where flexibility was proved only outside a subset of codimension 2.

Theorem 5.9. *The total coordinate space of a complete smooth T -variety of complexity one is flexible.*

Acknowledgments. We would like to thank Mikhail Zaidenberg for motivating questions and inspiring results and Ivan Arzhantsev for many useful remarks and suggestions. The first author started the project under Mobilnosc+ Polish Ministry of Science program, finished under DAAD PRIME program and was supported by the Foundation for Polish Science (FNP). The second author is grateful to the Dynasty Foundation for a partial financial support.

1. FLEXIBILITY OF AFFINE CONES

Lemma 1.1. *Let $X \subset \mathbb{A}^{n+1}$ be the affine cone over a projective variety $Y \subset \mathbb{P}^n$, smooth in codimension 1, which is not isomorphic to an affine space. Consider a subgroup $G \subset \operatorname{Aut} X$ such that*

- *the canonical $\mathbf{k}^*$ -action maps G -orbits to G -orbits and*
- *there is an orbit $Gx \subset X$, whose image under the projection $X \setminus \{0\} \rightarrow Y$ is the regular locus $Y_{\text{reg}} \subset Y$.*

Then $Gx = X_{\text{reg}}$.

Proof. As the action is equivariant, X_{reg} is a union of G -orbits, which projections coincide with Y_{reg} . Hence $X_{\text{reg}} = \bigcup_{\lambda \in \mathbb{G}_m} \lambda Gx$, where all G -orbits are closed in X_{reg} .

Let us show that there exists an open G -orbit $Gx = X_{\text{reg}}$. Assume the contrary. Then $\dim Gx = \dim Y$ and the stabilizer $S \subset \mathbb{G}_m$ of the orbit Gx is finite.

Denote by $X^\times$ the blow up of X at 0. It is the line bundle $\mathcal{O}_Y(-1)$ over Y . Consider a quotient morphism

$$\mu: X^\times \xrightarrow{/S} X' = X^\times / S$$

given by $t \mapsto t^{|S|}$ on every trivialization chart, where t is the coordinate of the fibers. Then $\mu(Gx)$ intersects each fiber at most once, so it is a meromorphic, nonvanishing section of the line bundle $X' \rightarrow Y$. But the subset $D \subset Y$, where the meromorphic section of the line bundle is not defined or zero, if non-empty, is a Cartier divisor, since it is defined locally by rational functions. On the other hand, $D \subseteq Y_{\text{sing}}$ is of codimension at least 2. Therefore, D is empty and $\mu(Gx)$ is a global section of X' , but X' is not a trivial bundle, as Y is not a point. A contradiction.

So, the group G acts on X_{reg} transitively. $\square$

Remark 1.2. We can weaken the second condition. Namely, if there is a G -orbit on X , whose image has complement of codimension ≥ 2 in Y , then this orbit is an open subset.

Lemma 1.3. *Let Y be an irreducible projective variety embedded into $\mathbb{P}^n$, and $U = Y \setminus \{x_0 = 0\}$ be a complement to a hyperplane section endowed with a $\mathbb{G}_a$ -action $\phi: \mathbb{G}_a \times U \rightarrow U$. Let $\pi: X \setminus \{x_0 = 0\} \rightarrow U$ be a natural projection, where $X = \text{AffCone } Y \subset \mathbb{A}^{n+1}$ is the affine cone over Y . Then X admits a homogeneous $\mathbb{G}_a$ -action $\hat{\phi}: \mathbb{G}_a \times X \rightarrow X$ such that*

- $\phi(\mathbb{G}_a \times \{\pi(x)\}) = \pi(\hat{\phi}(\mathbb{G}_a \times \{x\}))$ for any $x \in \pi^{-1}(U)$.
- $\hat{\phi}$ is trivial on $X \setminus \pi^{-1}(U)$.

In other words, $\pi: X \setminus \{x_0 = 0\} \rightarrow U$ provides an infinity-to-one correspondence between $\hat{\phi}$ -orbits and ϕ -orbits.

Proof. Let Y be defined by a homogeneous ideal $I \subset \mathbf{k}[x_0, \dots, x_n]$ which does not contain x_0 , $\mathbf{k}[X] = \mathbf{k}[x_0, \dots, x_n]/I$, and $U = \{x_0 \neq 0\} \subset Y$.

There exists a natural embedding $\rho: U \hookrightarrow X$, $\rho(U) = \{x_0 = 1\} \subset X$. On the other hand, $\pi: X \setminus \{x_0 = 0\} \rightarrow U$ is a trivial $\mathbb{G}_m$ -bundle. Therefore, we may extend the $\mathbb{G}_a$ -action ϕ on U to a $\mathbb{G}_a$ -action $\tilde{\phi}$ on $X \setminus \{x_0 = 0\}$ defined by a homogeneous locally nilpotent derivation $\tilde{\delta}$.

There exists $d \in \mathbb{N}$ such that $x_0^d \tilde{\delta}(x_i) \in \mathbf{k}[X]$ for $i = 1, \dots, n$. Since $x_0 \in \ker \tilde{\delta}$, a homogeneous derivation $\hat{\delta} = x_0^{d+1} \tilde{\delta}$ on X is locally nilpotent. The corresponding $\mathbb{G}_a$ -action $\hat{\phi}$ is homogeneous, coincides with ϕ on $U \cong \{x_0 = 1\} \subset X$, and is trivial on $\{x_0 = 0\} \subset X$. $\square$

Theorem 1.4. *Let Y be a projective variety covered by flexible normal open affine subsets U_i such that $Y \setminus U_i = D_i$, where D_i are effective divisors equivalent to a very ample divisor H . Then the affine cone $X = \text{AffCone}_H Y$ is flexible.*

Proof. The case $X = \mathbb{A}^{n+1}$ is trivial. We assume in sequel that the origin is a singular point of X .

For each element of the covering U_i there exists a finite number of $\mathbb{G}_a$ -actions $\{\phi_{ij}\}$ such that the orbit of the group generated by them is the regular locus of U_i . For each action ϕ_{ij} we can consider a lifted action $\hat{\phi}_{ij}$ on X as in Lemma 1.3. Let a subgroup $\{\hat{\phi}_{ij}\} \subset \text{SAut } X$ be generated by $\mathbb{G}_a$ -actions corresponding to all open subsets U_i . Then the image of the orbit of a regular point $Gx \subset X_{\text{reg}}$ under projection $X \setminus \{0\} \rightarrow Y$ is equal to Y_{reg} . Thus, the statement follows from Lemma 1.1, as the variety Y is normal, in particular smooth in codimension one. $\square$

Remark 1.5. We note that in Theorem 1.4 it is enough to assume that there exists an effective $\mathbb{Q}$ -divisor D' that has same support as D and is equivalent to H . Indeed, let $k > 0$ be such that kD' is a $(\mathbb{Z})$ -divisor, then we obtain $\mathbb{G}_a$ -actions on the affine cone over the Veronese embedding $v_k(Y)$. Following the proof of [KPZ11, Corollary 3.2] one can lift those actions to the cone over Y by [MM09, Theorem 3.1].

Remark 1.6. This criterion does not follow from [Per13, Thm 6]. Indeed, the $\mathbb{G}_a$ -actions on a flexible chart correspond to cylinders that are not necessarily H -polar.

Example 1.7. Consider $\mathbb{P}^n$ with coordinates $x_0, \dots, x_n$ and a smooth subvariety $Y = \mathbb{V}(x_0, f)$ of codimension 2, where f is a homogeneous polynomial of degree d . Let $q: X \rightarrow \mathbb{P}^n$ be the blowup in Y and $q': X' \rightarrow \mathbb{P}^n$ the combined blowup in Y and in the point $y = (1 : 0 : \dots : 0)$. We apply Theorem 1.4 to show that all affine cones over X and X' are flexible. Note that for $d \leq n$ we obtain Fano varieties via this construction.

Let $U_i := \mathbb{P}^n \setminus H_i := \mathbb{P}^n \setminus \mathbb{V}(x_i)$. The preimage $q^{-1}(U_0)$ is isomorphic to $U_0 \cong \mathbb{A}^n$, since Y does not intersect U_0 . For $i \neq 0$ the preimage $q^{-1}(U_i)$ is given by

$$V\left(\frac{f}{x_i^d} \cdot u - \frac{x_0}{x_i} \cdot v\right) \subset U_i \times \mathbb{P}^1.$$

Hence, $q^{-1}(U_i)$ is covered by the affine charts

$$U_i^0 := q^{-1}(U_i) \setminus [v = 0] \text{ and } U_i^\infty := q^{-1}(U_i) \setminus [u = 0],$$

the first one being an affine space and the second one being isomorphic to

$$V\left(\frac{f}{x_i^d} - \frac{x_0}{x_i} \cdot \frac{v}{u}\right) \subset \text{Spec } k\left[\frac{x_0}{x_i}, \dots, \frac{x_n}{x_i}, \frac{v}{u}\right].$$

In the notation of [AKZ12] this is a suspension over $\mathbb{A}^{n-1}$ and hence flexible by Theorem 0.2 loc. cit.

Given a divisor $D \subset \mathbb{P}^n$, we denote by $\tilde{D}$ its strict transform on the blowup. We also denote by E the exceptional divisor at Y and by H the pullback of the hyperplane. To see that the given covering is polar with respect to every ample divisor, note that

$$X \setminus U_0 = E \cup \tilde{H}_0; \quad X \setminus U_i^\infty = q^*H_i \cup \tilde{H}_0; \quad X \setminus U_i^0 = q^*H_i \cup \widetilde{\mathbb{V}(f)}.$$

Now, X has Picard group $\mathbb{Z}^2$ with generators $[H]$ and $[E]$. The effective cone is generated by $[E]$ and $[H] - [E]$ and the nef-cone by $[H]$ and $d[H] - [E]$. Moreover, $[\widetilde{\mathbb{V}(f)}] = d[H] - [E]$ and $[\tilde{H}_0] = [H] - [E]$ hold. So, for every of our affine charts we see that the components of the complement (1.7) span a cone in the Neron-Severi space containing the whole nef cone of X . Hence, every ample class can be expressed as a positive linear combination of the complement components.

Similarly, for X' we have the flexible charts

$$U_0^i := X \setminus (\tilde{H}_0 \cup E \cup \tilde{H}_i), \quad U_i^0 := X \setminus (\tilde{H}_i \cup \widetilde{\mathbb{V}(f)} \cup E'), \quad U_i^\infty := X \setminus (\tilde{H}_i \cup \tilde{H}_0 \cup E'),$$

the first two being affine spaces and the last one being a suspension as before. Here, we use the same notation as above and E' denotes the exceptional divisor of the blowup in the point. We see that the complements of the affine charts always consist of three components with classes corresponding to one of the triples

$$([H] - [E], [E], [H] - [E']), \quad ([H] - [E'], d[H] - [E], [E']), \quad ([H] - [E'], [H] - [E], [E']).$$

Further, each triple spans a cone containing the nef cone of X' , which is spanned by $d[H] - [E]$, $[H] - [E']$, and $[H]$. Hence, every ample class can be expressed as a positive linear combination of complement components.

2. SECANT OF SEGRE-VERONESE VARIETY

This section is based on [MOZ15], where the toric covering of a Segre variety was constructed. Here we generalize that construction to a Segre-Veronese variety and hence prove Theorem 2.21. Throughout this section by a *parameterization* of a variety Z we mean a dominant morphism from an open subset of an affine space to Z .

Definition 2.1. Given for each $i = 1, \dots, n$ a finite-dimensional vector space V_i and its symmetric power $S^{s_i}(V_i)$, $s_i \in \mathbb{Z}_{>0}$, the *Segre-Veronese variety*

$$X = v_{s_1}(\mathbb{P}(V_1)) \times \cdots \times v_{s_n}(\mathbb{P}(V_n)) \subset \mathbb{P}(S^{s_1}(V_1) \otimes \cdots \otimes S^{s_n}(V_n))$$

is defined as the embedding of the product $\mathbb{P}(V_1) \times \cdots \times \mathbb{P}(V_n)$ by the very ample line bundle $\mathcal{O}(s_1) \boxtimes \cdots \boxtimes \mathcal{O}(s_n)$.

We will be using an equivalent construction. Apart from (projective) Segre-Veronese varieties we will consider affine cones over them and refer to those as *Segre-Veronese cones*. They should not be confused with intersections of Segre-Veronese varieties with principal affine open subsets, which also play a crucial role.

For each V_i , $1 \leq i \leq n$, we denote $d_i = \dim V_i - 1$ and fix a basis $e_0^i, \dots, e_{d_i}^i$ of V_i . We also denote elements of the basis of $V_i^{\otimes s}$ by

$$e_{i_1, \dots, i_s}^i = e_{i_1}^i \otimes \cdots \otimes e_{i_s}^i.$$

Thus, the symmetric power is $S^{s_i}(V_i) =$

$$\{v \in V_i^{\otimes s} \mid (e_{i_1, \dots, i_s}^i)^*(v) = (e_{i_{\sigma(1)}, \dots, i_{\sigma(s)}}^i)^*(v) \text{ for any permutation } \sigma \in S_s\}.$$

This allows us to embed the Veronese cone into the Segre cone and obtain the following diagram:

$$\begin{array}{ccc} V_1 \oplus \cdots \oplus V_n & \xrightarrow{e} & V_1^{s_1} \oplus \cdots \oplus V_n^{s_n} \\ \downarrow \psi & & \downarrow \tilde{\psi} \\ S^{s_1}(V_1) \otimes \cdots \otimes S^{s_n}(V_n) & \subset & V_1^{\otimes s_1} \otimes \cdots \otimes V_n^{\otimes s_n}, \\ & & \swarrow s_{V_1} \otimes \cdots \otimes s_{V_n} \end{array}$$

where

- $e: (v_1, \dots, v_n) \mapsto (v_1, \dots, v_1, v_2, \dots, v_2, v_3, \dots, v_n)$ is the diagonal embedding,
- $\tilde{\psi}: (v_1^1, \dots, v_1^{s_1}, v_2^1, \dots, v_2^{s_2}, v_3^1, \dots, v_n^{s_n}) \mapsto v_1^1 \otimes \cdots \otimes v_n^{s_n}$ is the parameterization the Segre cone, which is a nonlinear map,
- $\psi = \tilde{\psi}|_{\text{im}(e)} \circ e$ is the parameterization of the Segre-Veronese cone, and
- $s_{V_i}: V_i^{\otimes s_i} \rightarrow S^{s_i}(V_i)$ are the natural symmetrizing projections.

Definition 2.2 ($\hat{\cdot}$, A , B).

- (i) For a vector space $V = V_{i_1} \otimes \cdots \otimes V_{i_k}$ we denote

$$\hat{V} := \{x \in V \mid (e_0^{i_1} \otimes \cdots \otimes e_0^{i_k})^*(x) = 1\}$$

and regard it as a vector space with basis $\{e_{j_1}^{i_1} \otimes \cdots \otimes e_{j_k}^{i_k} \mid j_1 + \cdots + j_k > 0\}$.

- (ii) We may, and often will, consider $\hat{V}$ as a complement to a hyperplane section of $\mathbb{P}(V)$ by the natural open embedding $\hat{V} \subset \mathbb{P}(V)$.
 (iii) We denote $A := V_1^{\otimes s_1} \otimes \cdots \otimes V_n^{\otimes s_n}$.
 (iv) We also define

$$B := \prod_{i=1}^n \hat{V}_i^{\times s_i} \subset V_1^{s_1} \oplus \cdots \oplus V_n^{s_n},$$

$$B' := \prod_{i=1}^n \hat{V}_i \subset V_1 \oplus \cdots \oplus V_n.$$

- (v) Finally, we denote $\pi: A \setminus \{0\} \rightarrow \mathbb{P}(A)$ and obtain the following diagram of open subsets:

$$\begin{array}{ccc} B' \subset V_1 \oplus \cdots \oplus V_n & \xrightarrow{e} & V_1^{s_1} \oplus \cdots \oplus V_n^{s_n} \supset B \\ \downarrow \psi & & \downarrow \tilde{\psi} \\ S^{s_1}(V_1) \otimes \cdots \otimes S^{s_n}(V_n) \subset V_1^{\otimes s_1} \otimes \cdots \otimes V_n^{\otimes s_n} \supset \hat{A} \\ & & \downarrow \pi \\ & & \mathbb{P}(A) \supset \hat{A}. \end{array}$$

Remark 2.3. Since $\mathbb{P}(S^{s_1}(V_1) \otimes \cdots \otimes S^{s_n}(V_n)) \subset \mathbb{P}(A)$, we can study the Segre–Veronese variety as a subvariety of $X = \pi \circ \tilde{\psi}(B') \subset \mathbb{P}(A)$. Note that the image of $\tilde{\psi}|_B$ does not contain the origin.

Cumulants. In this setting we may apply the (nonlinear) coordinate systems of B , called cumulants and presented in [MOZ15]. For the motivations to consider them, coming from algebraic statistics, we refer the reader to [SZ13, Zwi12, Zwi15]. A general mathematical setting for these methods is well presented in [CCM⁺14]. Further results are obtained for other varieties, e.g. Grassmannians and spinor varieties [MM15]. However, in other cases we do not obtain toric coverings. Still, we believe that similar methods can be applied to a larger class of secant and tangential varieties.

Definition 2.4 (indices). Basis elements of A are of form $e_{c_1^1, \dots, c_{s_1}^1}^1 \otimes \cdots \otimes e_{c_1^n, \dots, c_{s_n}^n}^n$ and are in natural correspondence with tuples $(c_1^1, \dots, c_{s_1}^1, \dots, c_1^n, \dots, c_{s_n}^n)$, where $0 \leq c_j^i \leq d_i$ for $1 \leq i \leq n$ and $1 \leq j \leq s_i$. Let us denote the set of these tuples by $C(A)$ and for each $c \in C(A)$ the corresponding basis element by $e(c)$. Finally, denote the dual basis elements by $x(c) = e(c)^*$. Similarly, we denote $C(\hat{A}) = C(A) \setminus \{(0, \dots, 0)\}$.

Definition 2.5 (degree, ordering). Given a tuple $c \in C(\hat{A})$, the number of its nonzero entries is called the *degree* of c . Given $c_1, c_2 \in C(\hat{A})$, we say that $c_1 \leq c_2$ if c_1 can be obtained from c_2 by setting some entries to zero.

Thus, we have a natural poset structure on $C(\hat{A})$, which induces a poset structure on the basis of $\hat{A}$. Simply speaking, the ordering on the basis is defined by replacing e_j^i by e_0^i in the tensor product elements.

Example 2.6. Consider $t_2 := e_1^1 \otimes e_0^1 \otimes e_5^2$, $t_1 := e_0^1 \otimes e_0^1 \otimes e_5^2$ and $t_0 := e_1^1 \otimes e_3^1 \otimes e_4^2$. We have $t_1 < t_2$ and t_1, t_2 are not comparable with t_0 .

Definition 2.7. Denote the set of indices

$$\text{Ind}(\hat{A}) = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \dots, \begin{bmatrix} 1 \\ s_1 \end{bmatrix}, \dots, \begin{bmatrix} n \\ 1 \end{bmatrix}, \dots, \begin{bmatrix} n \\ s_n \end{bmatrix} \right\}$$

and endow it with a natural lexicographic order, namely, the order of appearance above. Given $c \in C(\hat{A})$ and a subset of indices $I \subset \text{Ind}(\hat{A})$, we introduce the index tuple

$$c_I = (b_1^1, \dots, b_{s_n}^n), \quad \text{where } b_j^i = \begin{cases} c_j^i, & \begin{bmatrix} i \\ j \end{bmatrix} \in I, \\ 0, & \begin{bmatrix} i \\ j \end{bmatrix} \notin I. \end{cases}$$

Note that $\{b \in C(\hat{A}) \mid b \leq c\} = \{c_I \mid I \subset \text{Ind}(\hat{A})\}$. We will use either one-element subsets $I = \begin{bmatrix} i \\ j \end{bmatrix}$ or subsets of the following form, where $i_1, i_2 \in \text{Ind}(\hat{A})$:

$$I = [i_1 : i_2] := \{i \mid i_1 \leq i < i_2\} \subset \text{Ind}(\hat{A}).$$

Definition 2.8. A *thick interval partition* of a tuple $c \in C(\hat{A})$ of degree at least two is an increasing sequence of indices $\begin{bmatrix} 1 \\ 1 \end{bmatrix} = b_0 < \dots < b_k = \begin{bmatrix} n \\ s_n \end{bmatrix}$ such that $\deg c_{[b_i:b_{i+1}]} \geq 2$ for each i . The set of all thick interval partitions of c will be denoted by $IP(c)$. It is always nonempty as it contains $\left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} n \\ s_n \end{bmatrix} \right\}$.

Now we can recall the coordinate systems.

Definition 2.9 ([MOZ15, Sec. 2]). For each $c \in C(\hat{A})$ we denote

$$y(c) := \begin{cases} x(c), & \deg c = 1, \\ \sum_{(b) \leq (c)} (-1)^{\deg(c) - \deg(b)} x(b) \prod_{c_{j_0}^{i_0} \neq b_{j_0}^{i_0}} x(c_{[j_0]}^{i_0}), & \deg c > 1. \end{cases}$$

Then for each $c \in C(\hat{A})$ we introduce a function in $\mathbf{k}[\hat{A}]$

$$z(c) := \begin{cases} y(c), & \deg c = 1, \\ \sum_{(b_0, \dots, b_k) \in IP(c)} (-1)^k \prod_{m=1}^k y(c_{[b_{m-1}:b_m]}), & \deg c > 1. \end{cases}$$

Lemma 2.10. Each of sets $\{x(c)\}_{c \in C(\hat{A})}$, $\{y(c)\}_{c \in C(\hat{A})}$, and $\{z(c)\}_{c \in C(\hat{A})}$ is an algebraically independent system of functions generating $\mathcal{O}(\hat{A})$. In other words, $\{z(c)\}$ is a coordinate system on $\hat{A}$ as an affine space.

Proof. Since $y(c)$ is a sum of $x(c)$ and of terms of smaller degree, the endomorphism of $\mathbf{k}[\hat{A}]$ that maps $x(c)$ to $y(c)$ for each c is invertible. Same holds for $\{y(c)\}$ and $\{z(c)\}$. So, the statement follows. $\square$

Secant. The secant variety $\text{Sec } X \subset \mathbb{P}(A)$ of the Segre–Veronese variety X is parameterized by a map

$$(1) \quad \text{sec}_X: \mathbb{A}^1 \times B' \times B' \rightarrow \hat{A}, \quad (t, v, w) \mapsto \pi(t \cdot \psi(v) + (1-t) \cdot \psi(w)).$$

Hereinafter, given a tuple of degree one, we denote the index of its only non-zero entry by $\begin{bmatrix} i_\circ \\ j_\circ \end{bmatrix}$. Generalizing [MOZ15, Lemma 3.1], we obtain the following result.

Lemma 2.11. Let $\{(e_j^i)^{i*} \mid 1 \leq i \leq n, 0 \leq j \leq d_i\}$ be the set of coordinate functions on the first copy of B' in (1) and $\{(e_j^i)^{i**}\}$ the respective set on the second one.* Then

$$\text{sec}_X^*: z(c) \mapsto \begin{cases} t(e_{c_{j_\circ}^{i_\circ}}^{i_\circ})^{i*} + (1-t)(e_{c_{j_\circ}^{i_\circ}}^{i_\circ})^{i**}, & \deg c = 1 \text{ with } c_{j_\circ}^{i_\circ} \neq 0, \\ t(1-t)(1-2t)^{\deg(z(c))-2} \prod_{c_j^i \neq 0} ((e_{c_j^i}^i)^{i*} - (e_{c_j^i}^i)^{i**}), & \deg c > 1, \end{cases}$$

for each $c \in C(\hat{A})$.

Proof. Let Y be the cone over the Segre product $\mathbb{P}(V_1)^{s_1} \times \dots \times \mathbb{P}(V_n)^{s_n}$. Then the secant of Y is parameterized by

$$\text{sec}_Y: \mathbb{A}^1 \times B \times B \rightarrow \hat{A}, \quad (t, v, w) \mapsto \pi(t\tilde{\psi}(v) + (1-t)\tilde{\psi}(w)).$$

Thus, $\text{sec}_X = \text{sec}_Y \circ (\text{id} \times e \times e)$. The statement follows after applying [MOZ15, Lemma 3.1] to sec_Y . $\square$

*That is, $(e_j^i)^{i*}(v) = v_j^i$ and $(e_j^i)^{i**}(w) = w_j^i$ respectively.

Torus action. We can infer the following decomposition of sec_X .

Definition 2.12. Let us introduce a reparameterization $\text{rep}: \mathbb{A}^1 \times B' \times B' \rightarrow \mathbb{A}^1 \times B' \times B'$,

$$\text{rep}: (t, v, w) \mapsto \left(\frac{t(1-t)}{(1-2t)^2}, tv + (1-t)w, (1-2t)(w-v) \right)$$

and a *monomial* map $m: \mathbb{A}^1 \times B' \times B' \rightarrow \hat{A}$,

$$m^*: z(c) \mapsto \begin{cases} (e_{c_{j_o}^{i_o}}^{i_o})'^*, & \deg c = 1 \text{ with } c_{j_o}^{i_o} \neq 0, \\ t \prod_{c_j^i \neq 0} (e_{c_j^i}^i)''^*, & \deg c > 1. \end{cases}$$

Lemma 2.13. *There is a decomposition $\text{sec}_X = m \circ \text{rep}$. In particular, m is a monomial parameterization of $\text{Sec } X$.*

Proof. Straightforward. $\square$

This monomial parameterization of $\text{Sec } X$ already provides a structure of a toric variety on $\text{im}(m) = \hat{A} \cap \text{Sec } X$, hence provides us with a toric chart of $\text{Sec } X$. Below we describe in detail its structure.

Definition 2.14. Let us introduce the following closed subsets of $\hat{A}$:

$$\begin{aligned} \hat{A}_1 &= \{z(c) = 0 \mid \deg c > 1\}, \\ \hat{A}_2 &= \{z(c) = 0 \mid \deg c = 1\}, \\ S_2 &= \text{Sec } X \cap A_2. \end{aligned}$$

Definition 2.15 (P). Consider the lattice $M = \bigoplus_{\substack{1 \leq i \leq n \\ 1 \leq j \leq d_i}} \mathbb{Z} \chi_j^i$, where $\chi_j^i = (e_j^i)^*$.

We introduce the lattice polytope $P \subset M \otimes \mathbb{Q}$ defined by inequalities

$$\begin{cases} (\chi_j^i)^* \geq 0, & 1 \leq i \leq n, 1 \leq j \leq d_i, \\ \sum_{j=1}^{d_i} (\chi_j^i)^* \leq s_i, & 1 \leq i \leq n, \\ \sum_{\substack{1 \leq i \leq n \\ 1 \leq j \leq d_i}} (\chi_j^i)^* \geq 2. \end{cases}$$

Proposition 2.16. *In the terminology above,*

- (i) $\text{Sec } X \cap \hat{A} = \hat{A}_1 \times S_2$ via natural projections along coordinates $z(c)$,
- (ii) $\hat{A}_1 = X \cap \hat{A} \cong \mathbb{A}^N$, where $N = \sum_{i=1}^n \dim \hat{V}_i$,
- (iii) $S_2 \cong \text{AffCone}(X_P)$, where X_P is a projective toric variety with polarization corresponding to the polytope P , e.g. see [CLS11, §2.3] for the construction.

Proof. The morphism m is a direct product of $m|_{\{0\} \times B' \times \{0\}}$ and $m|_{\mathbb{A}^1 \times \{0\} \times B'}$, which respectively parameterize $\hat{A}_1$ and S_2 . This implies (i). The Segre–Veronese variety X is also parameterized by $m|_{\{0\} \times B' \times \{0\}}$, thus (ii) holds.

Let us consider the standard torus $T = \text{Spec } \mathbf{k}[M] \subset B'$ and a projectivization $\pi_z: \hat{A} \setminus \{0\} \rightarrow \mathbb{P}(\hat{A})$ along $z(c)$ -coordinates. Then $X_P = \overline{\pi_z(S_2)}$ is parameterized by a T -equivariant map $\pi_z \circ m|_{\{0\} \times \{0\} \times B'}$, which is defined by the set of monomials $\{(e_{c_j^i}^i)^* \mid \deg c > 1\} \subset \mathcal{O}(B')$ corresponding exactly to lattice points $P \cap M$. This implies (iii). $\square$

Proposition 2.17. *Let $\text{Tan } X \subset \mathbb{P}(A)$ be the tangential variety of X . Then*

$$\text{Tan } X \cap \hat{A} = \hat{A}_1 \times X'_P,$$

where $X'_P \subset \hat{A}_2$ is a nondegenerate toric variety parameterized by $m|_{\{-\frac{1}{4}\} \times \{0\} \times B'}$.

Proof. We present here a sketch of a proof for $\mathbf{k} = \mathbb{C}$, see [MOZ15] and [MM15, Lem. 3.3] for a complete proof. Consider $(\epsilon^{-1}, v, v + \epsilon w) \in \mathbb{A}^1 \times B' \times B'$. If $\epsilon \rightarrow 0$, then $\sec_X(\epsilon^{-1}, v, v + \epsilon w)$ tends to an element of $T_{\pi \circ \psi(v)} X \hookrightarrow \text{Tan } X$ corresponding to w . On the other hand, by Lemma 2.11,

$$\lim_{\epsilon \rightarrow 0} z(c)(\sec_X(\epsilon^{-1}, v, v + \epsilon w)) \rightarrow \begin{cases} v_{c_{j_0}^{i_0}}^{i_0} - w_{c_{j_0}^{i_0}}^{i_0}, & \deg c = 1 \text{ with } c_{j_0}^{i_0} \neq 0, \\ -\frac{1}{4} \prod_{c_j^i \neq 0} 2w_{c_j^i}^i, & \deg c > 1. \end{cases}$$

Thus, the decomposition follows. It remains to check that X'_P is nondegenerate. Indeed, $\mathcal{O}(X'_P) \hookrightarrow \mathcal{O}(B')$ does not contain invertible elements. $\square$

These propositions imply the following relationship of the tangential and secant varieties, which, in its turn, implies Zak's theorem [Zak05] for Segre–Veronese varieties.

Corollary 2.18. *The following conditions are equivalent:*

- (i) P is not contained in the hyperplane $\sum_{i,j} (\chi_j^i)^* = 2$,
- (ii) $\dim \text{Sec } X = 2 \dim X + 1$,
- (iii) $\text{Sec } X \neq \text{Tan } X$,
- (iv) $\dim \text{Tan } X = \dim \text{Sec } X - 1$.

Then $\text{Sec } X$ is called non-degenerate.

Proof. Let $d = \dim X$. As a toric variety, X is represented by a d -dimensional polytope S , which is a product of simplices. Then P is the intersection of S with the halfspace $\sum_{i,j} (\chi_j^i)^* \geq 2$.

$i) \Rightarrow ii)$: The assumption implies $\dim P = d$. Hence, $\dim S_2 = d + 1$ and $\dim \text{Sec } X = \dim \hat{A}_1 + \dim S_2 = 2d + 1$.

$i) \Rightarrow iv)$: As before, $\dim \text{Cone}(P) = d$. Hence, $\dim \text{Tan } X = \dim \hat{A}_1 + d = 2d$.

The implications $ii) \Rightarrow iii)$ and $iv) \Rightarrow iii)$ are obvious.

$iii) \Rightarrow i)$: If P is contained in the hyperplane, then all monomials corresponding to lattice points in P are of same degree. In particular, $X'_P \cong \text{AffCone}(X_P)$. $\square$

Example 2.19. 1) Consider the Veronese surface $Y = \mathbb{P}^2 \hookrightarrow \mathbb{P}^5$. It is a projective toric variety corresponding to the simplex $S = \text{conv}(0, 2\chi_1, 2\chi_2) \subset \langle \chi_1, \chi_2 \rangle \cong \mathbb{Z}^2$, i.e. parameterized by characters of a two-dimensional torus that correspond to the lattice points of S . By Proposition 2.16, the variety X_P that defines the secant is parameterized by $P = S \cap \{\chi_1^* + \chi_2^* \geq 2\}$, i.e., by lattice points $2\chi_1, \chi_1 + \chi_2, 2\chi_2$. So, both factors of $\text{Sec } X \cap \hat{A} = \hat{A}_1 \times S_2$ are of dimension two, so $\dim \text{Sec } X = 4$. Thus, the secant variety is degenerate and defined by $z(1, 2)^2 = z(1, 1)z(2, 2)$.

2) Consider the Segre product $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. The representing polytope is a cube $[0, 1]^3$. The secant variety is represented by a polytope with vertices $(1, 1, 0), (1, 0, 1), (0, 1, 1), (1, 1, 1)$. We see that the affine cone over it is the whole affine space; indeed in this case the secant variety is non-degenerate and fills the whole ambient space. The tangential variety is a hypersurface defined by the equation $z(1, 1, 0)z(1, 0, 1)z(0, 1, 1) = z(1, 1, 1)^2$.

Theorem 2.20. *The tangential and secant varieties of a Segre–Veronese variety are covered by complements of hyperplane section. Each complement is an affine toric variety without torus factors. In case of the secant variety, these are always normal toric varieties. In case of the tangential variety they are normal if the underlying variety is the Segre product.*

Proof. By [Ver09, Theorem 2.2] we know that $\text{Sec } X$ is normal. By [MOZ15, Proposition 8.5] we know that $\text{Tan } X$ is normal, when X is the Segre product. The open subsets $\text{Sec } X \cap \hat{A}$ and $\text{Tan } X \cap \hat{A}$ are toric varieties by Propositions 2.16 and 2.17.

Moreover, they do not contain torus factors, since they are products of an affine space $\hat{A}_1$ with either an affine cone over a projective toric variety X_P or a non-degenerate toric variety X'_P . By taking such subsets for various choices of basis vectors e_0^i , $i = 1, \dots, n$, we obtain the statement. $\square$

Theorem 2.21. *Let $X = v_{s_1}(\mathbb{P}(V_1)) \times \dots \times v_{s_n}(\mathbb{P}(V_n))$ be a Segre–Veronese variety. Then the affine cone over the secant variety of X is flexible. Further, if $s_1 = \dots = s_n = 1$, then also the affine cone over the tangential variety of X is flexible.*

Proof. By [AKZ12, Theorem 0.2] we know that all the affine charts from Theorem 2.20 are flexible. Now, by Theorem 1.4 we obtain flexibility of the cone. $\square$

Example 2.22. Consider the third Veronese embedding $X = v_3(\mathbb{P}^1)$, it is represented by characters in the interval $S = \text{conv}(0, 3)$. Then $\text{Tan } X \cap \hat{A} \cong \mathbb{A}^1 \times X'_P$, where the monoid of characters associated to the toric variety X'_P is generated by $\{2, 3\}$. Namely, X'_P is the curve with a cusp singularity at the origin. Thus, $\text{Tan } X$ is a two-dimensional surface, whose singular locus is the curve X . This example can be generalized to tangential varieties of other Segre–Veronese varieties provided that at least one of the Veronese factors is of degree at least 3.

3. THE COMBINATORIAL DESCRIPTION OF T-VARIETIES

We consider a normal variety X with an effective action of an algebraic torus $T \cong (\mathbb{G}_m)^r$. Then X is called a T -variety of complexity $(\dim X - \dim T)$. In the following we restrict ourselves to the case of rational T -varieties of complexity one. Our general reference is [AIP⁺11].

Let us denote the character lattice of the torus T by M and the dual lattice by N . For the associated vector spaces we write $M_{\mathbb{Q}}$ and $N_{\mathbb{Q}}$.

For a polyhedron $\Delta \subset N_{\mathbb{Q}}$ we consider its tail cone $\text{tail}(\Delta) := \{v \in N_{\mathbb{Q}} \mid \Delta + \mathbb{Q}_{\geq 0} \cdot v = \Delta\}$. Now, we consider polyhedral complexes Ξ in $N_{\mathbb{Q}}$ such that the set of tail cones has the structure of a fan, which is called the tail fan of Ξ and will be denoted by $\text{tail}(\Xi)$. Consider a pair $\mathcal{S} = (\sum_{P \in \mathbb{P}^1} \mathcal{S}_P \otimes P, \deg \mathcal{S})$ where $\mathcal{S}_P$ are polyhedral complexes in $N_{\mathbb{Q}}$ with some common tail fan Σ and $\deg \mathcal{S} \subset |\Sigma|$. Here, $\sum_P \mathcal{S}_P \otimes P$ is just a formal sum. The complexes $\mathcal{S}_P$ are called *slices* of $\mathcal{S}$. We assume that there are only finitely many slices that differ from the tail fan $\text{tail}(\mathcal{S}) := \Sigma$. The set of the points $P \in \mathbb{P}^1$ such that $\mathcal{S}_P \neq \Sigma$ is called the support of $\mathcal{S}$ and will be denoted by $\text{supp } \mathcal{S}$. Note that for every full-dimensional $\sigma \in \Sigma$ there is a unique polyhedron Δ_P^σ in $\mathcal{S}_P$ with $\text{tail}(\Delta_P^\sigma) = \sigma$.

Definition 3.1 (f-divisor). A pair $\mathcal{S}$ as above is called an *f-divisor* if for any full-dimensional $\sigma \in \text{tail}(\mathcal{S})$ we have either $\deg \mathcal{S} \cap \sigma = \emptyset$ or

$$\sum_P \Delta_P^\sigma = \deg \mathcal{S} \cap \sigma \subsetneq \sigma.$$

An f-divisor as above corresponds to a rational T -variety of complexity one, see [IS11, Section 1]. Moreover, this correspondence is even functorial. In particular, invariant open subvarieties correspond to f-divisors $\mathcal{S}'$, such that $\mathcal{S}'_P \subset \mathcal{S}_P$ as sets of polyhedra and $\deg \mathcal{S}' = |\text{tail}(\mathcal{S}')| \cap \deg \mathcal{S}$. For simplicity we write $\mathcal{S}' \subset \mathcal{S}$ in this situation. As a consequence of Proposition 1.6 in [IS11], f-divisors $\mathcal{S}_1, \dots, \mathcal{S}_\ell \subset \mathcal{S}$ give rise to an open covering if and only if their slices cover the slices of $\mathcal{S}$, i.e.

$$|\mathcal{S}| = \bigcup_i |\mathcal{S}_i|.$$

Remark 3.2. Affine charts correspond to f-divisors $\mathcal{S}$ such that $\mathcal{S}_P$ consists of a single polyhedron (and its faces) and $\deg \mathcal{S} = \sum_{P \in \mathbb{P}^1} \mathcal{S}_P$. These objects are called p-divisors in [IS11, AIP⁺11].

Example 3.3. In Figure 1 we sketched the non-trivial slices of an f-divisor as well as its degree. It describes the blowup of the quadric threefold in one point, see [Süß14].




FIGURE 1. An f-divisor

Lemma 3.4 ([IV12, Remark 1.8.]). *An f-divisor describes a subtorus action on a toric variety if and only if $\mathcal{S}_P$ equals a lattice translate of the tail fan for all but at most two $P \in \mathbb{P}^1$.*

In the language of f-divisors we also may describe torus invariant Cartier divisors by *support functions*. A support function h on a polyhedral subdivision Ξ is a continuous function that is affine linear on every polyhedra in Ξ . We denote by $\text{lin } h$ the linear part of h . This is a piecewise linear function on the tail fan defined as follows:

$$(\text{lin } h)(v) := h(w + v) - h(w)$$

for some $w \in \Delta \in \Xi$ with $v \in \text{tail}(\Delta)$.

Definition 3.5 (Support function on $\mathcal{S}$). A *support function* h on an f-divisor $\mathcal{S}$ is a collection $\{h_P\}_{P \in \mathbb{P}^1}$ of support functions on $\mathcal{S}_P$ such that

- (i) all h_P have the same linear part, which will be denoted by $\text{lin } h$,
- (ii) only finitely many of them differ from $\text{lin } h$.

We have two kinds of torus invariant prime divisors on $X(\mathcal{S})$. *Horizontal* prime divisors correspond to rays $\rho \in \text{tail}(\mathcal{S})^{(1)}$ that do not intersect $\deg \mathcal{S}$ and are denoted by D_ρ . *Vertical* prime divisors correspond to vertices v in the subdivisions $\mathcal{S}_P$ and are denoted by $D_{P,v}$. Now, the divisor corresponding to the support function h is given by

$$D_h = - \sum_{\rho} (\text{lin } h)(\rho) \cdot D_\rho - \sum_{P,v} \mu(v) \cdot h_P(v) D_{P,v},$$

where we identify the ray with the ray generator and $\mu(v)$ denotes the minimal positive integer such that $\mu(v) \cdot v$ is a lattice element. In particular,

- (i) if $h_P \leq 0$ for all $P \in \mathbb{P}^1$, then D_h is effective.
- (ii) in this case $X(\mathcal{S}) \setminus \text{supp } D_h$ is given by the f-divisor

$$[h = 0] := \left(\sum_P [h_P = 0] \otimes P, \quad \deg \mathcal{S} \cap [\text{lin } h = 0] \right),$$

where $[h_P = 0]$ denotes the polyhedral subcomplex of $\mathcal{S}_P$ consisting of those polyhedra on which h_P vanishes.

By [PS11], every invariant Cartier divisor arises in this way. We have $D_h \sim D_{h'}$ if and only if $h_P - h'_P$ is affine linear for every P , i.e. $h_P - h'_P = \langle u, \cdot \rangle + a_P$, and

$\sum_P a_P = 0$. Moreover, we have a criterion for ampleness expressed in the following notation. We denote by $\square_h$ the polytope given by

$$(2) \quad \square_h = \{u \in M_{\mathbb{Q}} \mid \langle u, \cdot \rangle \geq (\text{lin } h)\}$$

and consider concave piecewise affine function h_P^* on $\square_h$ as “dual” of h_P :

$$h_P^*(u) := \inf_v (\langle u, v \rangle - h_P(v)).$$

The definition implies that $h_P^*(u)$ is finite for $u \in \square_h$.

Theorem 3.6. *If D_h is ample, then h_P is strongly concave for every $P \in \mathbb{P}^1$ and $h_P^*(u) \geq 0$ for every $u \in \square_h$.*

Proof. By [PS11, Theorem 3.28], h_P has to be strongly concave and by [IS11, Prop. 3.1(i)] we get that $h_P^*(u) \geq 0$. $\square$

4. CONES OVER PROJECTIVE T-VARIETIES

From now on we assume that the T -varieties we consider are proper over the base, i.e. that all slices $\mathcal{S}_P$ are *subdivisions* of $N_{\mathbb{Q}}$.

Definition 4.1 (Equivariant covering by toric charts). A T -variety is called *equivariantly covered by toric charts*, if there is an open covering by toric varieties U_i such that the torus T acts as a subtorus of the embedded torus of U_i .

Lemma 4.2. *The T -variety $X(\mathcal{S})$ is equivariantly covered by toric charts if and only if for every maximal polyhedron Δ in $\mathcal{S}_P$, $P \in \mathbb{P}$, all but at most two slices contain a lattice translate of $\text{tail}(\Delta)$. In particular, either $X(\mathcal{S})$ itself is toric or there is at most one $P \in \mathbb{P}^1$ such that $\mathcal{S}_P$ does not contain a lattice vertex.*

Proof. The first part is a corollary of Lemma 3.4. To prove the last statement, we consider two points P, Q such that $\mathcal{S}_P, \mathcal{S}_Q$ contain only non-lattice vertices. Now, consider a third point R and a maximal polyhedron $\Delta \subset \mathcal{S}_R$. Since there is no lattice translate of $\text{tail}(\Delta)$ in $\mathcal{S}_P$ and $\mathcal{S}_Q$, Δ itself must be a translated cone. Notice that all maximal cones in $\mathcal{S}_R$ must be translated by the same lattice point. Indeed, otherwise they would not cover $N_{\mathbb{Q}}$ and there would exist a different maximal dimensional polyhedron that could not be a lattice translate of its tail cone. Hence, for any $R \notin \{P, Q\}$ the slice $\mathcal{S}_R$ is just a translated tail fan. By Lemma 3.4, $X(\mathcal{S})$ is a toric variety. $\square$

Remark 4.3. By [APS14, Appendix] this criterion is fulfilled for all smooth rational T -varieties of complexity one. Hence, they are covered by affine spaces.

To get flexibility for every affine cone we need to strengthen the condition in Lemma 4.2.

Theorem 4.4. *Let $X = \mathfrak{X}(\mathcal{S})$ be a T -variety such that for any maximal polyhedron $\Delta \in \mathcal{S}_y$, $y \in \mathbb{P}^1$, at most two slices contain a polyhedron with the same tail cone $\text{tail}(\Delta)$ that is not a lattice translate of $\text{tail}(\Delta)$. Then for every very ample divisor H the corresponding affine cone is flexible.*

Proof. By Theorem 1.4 it is enough to show that there exists an equivariant H -polar covering by toric charts. Let us first rephrase this condition in terms of f -divisors. For every maximal polyhedron $\Delta_P \in \mathcal{S}_P$ there exists a strictly concave non-positive support function h on $\mathcal{S}$ corresponding to an effective divisor $D_h \sim H$, such that $[h_P = 0] = \Delta_P$ (i.e. $h_P|_{\Delta_P} \equiv 0$ and negative elsewhere). For being a toric covering additionally we have to impose that $[h = 0]$ has only two slices that are not lattice translates of the tail fan.

We now construct such a covering for some very ample divisor $H = D_h$ corresponding to a support function h . Fix a maximal polyhedron $\Delta \subset \mathcal{S}_Q$. Then $h_Q|_\Delta$ is affine linear, i.e. $h_Q(v) = \langle u, v \rangle + a$. By concavity this implies $u \in \square_h$, with $\square_h$ defined as in (2). We now consider $h' := h - u$ with $h'_P(v) := h_P(v) - \langle u, v \rangle$. Now, h'_P is again strongly concave and obtains its maximum at a polyhedron Δ_P with tail cone $\text{tail}(\Delta_P) = \text{tail}(\Delta)$. Moreover, by construction we have $0 \in \square_{h'}$.

By our precondition, we may assume without loss of generality that for every point $R \in \mathbb{P} \setminus \{0, \infty\}$ the polyhedron Δ_R is a lattice translate of $\text{tail}(\Delta)$. Assume further $Q \neq \infty$ and introduce h^∞ by

$$\begin{cases} h_P^\infty(v) := h'_P(v) - \max \text{Im } h'_P & \text{for } P \neq \infty, \\ h_\infty^\infty(v) := h'_\infty(v) + \sum_{P \neq \infty} \max \text{Im } h'_P. \end{cases}$$

It remains to check that $h'_\infty(v) + \sum_{P \neq \infty} \max \text{Im } h'_P \leq 0$ to see that D_{h^∞} is indeed effective. Recall that we have $0 \in \square_{h'}$. The claim follows from the ampleness of $D_{h'}$ and Theorem 3.6.

Now, by construction we have $D_{h^\infty} \sim H$ and $[h_P^\infty = 0] = \Delta$. Moreover, $[h_P^\infty = 0]$ is a lattice translate of $\text{tail}(\Delta)$ for each $P \notin \{0, \infty\}$. Then it describes a toric chart.

Taking these toric charts for every maximal polyhedron provides us with an H -polar covering. Now, our result follows by Theorem 1.4. $\square$

Theorem 4.5. *All affine cones over the Fano threefolds Q , 2.29, 2.30, 2.31, 2.32, 3.8, 3.18, 3.19, 3.23, 3.24, 4.4, and elements of the families 2.24, 3.10 admitting a 2-torus action in Mori–Mukai’s classification are flexible.*

Proof. For all Fano threefolds from Theorem 4.5 the corresponding f-divisors are listed in [Süß14]. One can easily check that the precondition of Theorem 4.4 is fulfilled in every case. $\square$

Example 4.6. Let us illustrate the difference of assumptions in Lemma 4.2 and Theorem 4.4. In the lemma we are allowed to *choose* the polyhedron with the given tail cone. Hence, if we consider the variety given by the slice $(-\infty, -1], [-1, 1], [1, \infty)$ taken three times, then it does satisfy the assumptions. Indeed, if we take the maximal polytope $[-1, 1]$ in one slice, in other two slices we can take just the vertex $\{1\}$, which is a lattice shift of the tail cone $\{0\}$. On the other hand in the theorem we ask for *all* polyhedra with the given tail cone. Here, we get three times $[-1, 1]$ which is not a lattice translate of $\{0\}$. Such a difference is only possible for cones that are not full-dimensional.

Example 4.7. We are coming back to the blowup of the quadric threefold from Example 3.3. We may check that the corresponding f-divisor in Figure 1 fulfills the condition of Theorem 4.4. Hence, all affine cones over the blowup of the quadric threefold are flexible.

Example 4.8. The hypersurface $\mathbb{V}(x_0y_0^2 + x_1y_1^2 + x_2y_2^2) \subset \mathbb{P}^2 \times \mathbb{P}^2$ is 2.24 from our list in Theorem 4.5. Hence, every affine cone over this variety is flexible. In particular, this is true for the cone over the Segre embedding.

5. TOTAL COORDINATE SPACES

We recall the definition of Cox rings.

Definition 5.1 (Cox sheaf, Cox ring, universal torsor, total coordinate space). Let X be a normal variety, whose class group is a free abelian group generated by divisor classes $D_1, \dots, D_r$. The *Cox sheaf* of X is defined by

$$\mathcal{R} = \bigoplus_{\vec{a} \in \mathbb{Z}_{\geq 0}^r} \mathcal{O}(\sum_{i=1}^r a_i D_i).$$

It becomes a sheaf of $\mathcal{O}_X$ -algebras via the usual multiplication of sections. Its global sections are called the *Cox ring* denoted by $\mathcal{R}(X)$.

The relative spectrum $\hat{X} = \text{Spec}_X(\mathcal{R})$ is called the *universal torsor* of X . It is an open subset of the absolute spectrum $\overline{X} = \text{Spec}(\mathcal{R}(X))$, which is called the *total coordinate space* of X . By construction, the Cox ring is graded by the class group of X inducing an action of the torus $\text{Spec } k[\text{Cl}(X)]$ on the total coordinate space.

In the following we are studying flexibility of total coordinate spaces for several classes of varieties.

5.1. Del Pezzo surfaces. Since the smooth del Pezzo surfaces of degrees 6, 7, 8, and 9 are toric, their total coordinate spaces are just affine spaces and hence flexible. The remaining del Pezzo surfaces are blowups X_r of $\mathbb{P}^2$ in r points of general position, where $4 \leq r \leq 8$. Their Cox rings are described for example in [BP04], [Der07], and [TVV09].

An exceptional curve on X is a curve of self-intersection -1 and anti-canonical degree 1. On every del Pezzo surface there are only finitely many of them. We will use the following facts from [BP04].

Theorem 5.2 ([BP04, Thm 3.2 & Prop. 3.4]). *Let $e_1, \dots, e_N$ be sections corresponding to exceptional curves on a del Pezzo surface X_r and I be the ideal of their relations. Then*

- (i) $\mathcal{R}(X_r) = \mathbf{k}[e_1, \dots, e_N]/I$ for $4 \leq r \leq 7$;
- (ii) $\mathcal{R}(X_8) = \mathbf{k}[e_1, \dots, e_N]/I \oplus \langle f_1, f_2 \rangle_{\mathbf{k}}$ as a vector space,
where $f_1, f_2 \in H^0(X_8, \mathcal{O}(-K_{X_8}))$ are elements of degree one with respect to the $\mathbb{Z}$ -grading by the anti-canonical degree of a divisor class.

Theorem 5.2 shows that the Cox ring is generated by elements of degree 1 and $Y_r := \text{Proj}(\mathcal{R}(X_r))$ comes with an embedding into $\mathbb{P}^{N-1}$, where N (resp. $N-2$) is the number of exceptional curves in the case $4 \leq r \leq 7$ (resp. $r = 8$).

In this situation the total coordinate space $\overline{X}_r$ is the affine cone over this embedding.

Proposition 5.3. *Let e be a section corresponding to an exceptional curve. Then the principal open subset $(Y_r)_e$ is isomorphic to X_{r-1} .*

Proof. This can be found for example in the proof of Proposition 3.4 in [BP04]. $\square$

Theorem 5.4. *The total coordinate spaces of smooth del Pezzo surfaces are flexible.*

Proof. As said above, it is enough to check X_r with $4 \leq r \leq 8$. We will go by induction. The del Pezzo surface X_3 is toric. Therefore it has a flexible total coordinate space. Now, consider X_r with $4 \leq r \leq 8$. Then we have seen that $\overline{X}_r$ is the affine cone over Y_r . Moreover, principal open subsets corresponding to sections of exceptional curves are isomorphic to $\overline{X}_{r-1}$ and hence flexible by induction hypothesis. It remains to check that these principal open subsets cover Y_r to conclude flexibility of $\overline{X}_r$ from Theorem 1.4. For $4 \leq r \leq 7$ this follows directly from Theorem 5.2 (i). For the case $r = 8$ we have to take care for the remaining generators. By Theorem 5.2 (ii) their squares are contained in the ideal $(e_1, \dots, e_N)$ generated by the sections corresponding to exceptional curves, but then the common vanishing of $e_1, \dots, e_N$ implies the vanishing of the remaining generators and hence $Y_r = \bigcup_{i=1}^N (Y_r)_{e_i}$. $\square$

5.2. Smooth complexity-one T-varieties. In [HS10] the Cox rings of T-varieties are studied. For the case of complexity-one action they have a very particular form.

Proposition 5.5 ([HS10, Corollary 4.9]). *Let $\mathcal{S}$ be an f -divisor and let us denote by $\mathcal{S}^\times$ the subset of rays in $\text{tail}(\mathcal{S})^{(1)}$ that do not intersect $\deg \mathcal{S}$. Then the Cox ring of $X(\mathcal{S})$ is given by*

$$\frac{\mathbf{k}[S_\rho, T_{P,v} \mid \rho \in \mathcal{S}^\times, P \in \text{supp } \mathcal{S}, v \in \mathcal{S}_P^{(0)}]}{\langle z \cdot T^{\mu(0)} + T^{\mu(\infty)} + T^{\mu(z)} \mid z \in \text{supp } \mathcal{S} \cap \mathbf{k}^* \rangle}$$

where $T^{\mu(P)} := \prod_{v \in \mathcal{S}_P^{(0)}} T_{P,v}^{\mu(v)}$ and $\mu(v)$ denotes the minimal positive integer such that $\mu(v) \cdot v$ is a lattice element.

If we impose the additional condition that the T-variety is equivariantly covered by toric charts (which is fulfilled in the smooth case), then we can conclude the following.

Proposition 5.6. *The Cox ring of a complexity-one T-variety equivariantly covered by toric charts is isomorphic to*

$$\mathbf{k}[S_1, \dots, S_{n_S}; T_{\ell,j} \mid 0 \leq \ell \leq m, 1 \leq j \leq n_\ell] / \langle z_\ell \cdot A_0 + A_1 + A_\ell \mid 2 \leq \ell \leq m \rangle,$$

where

- (i) $N, m, n_0, \dots, n_m \in \mathbb{Z}_{>0}$;
- (ii) $z_2, \dots, z_m$ are distinct elements of $\mathbf{k}^*$;
- (iii) for $\ell = 0, \dots, m$, A_ℓ is a monomial in $\mathbf{k}[T_{\ell,1}, \dots, T_{\ell,n_\ell}]$;
- (iv) for $\ell = 1, \dots, m$ the monomial A_ℓ is linear in at least one variable.

Moreover, if X is Fano, then we may assume that A_ℓ for $\ell \geq 3$ is linear in each variable.

Proof. The first statements follow directly from Proposition 5.5 and Lemma 4.2. For the Fano case note that the Cox ring of a Fano variety is log-terminal by [Bro13, GOST15] and factorial by [BH03]. Hence, by Remark 6.4 in [LS13] we obtain the last statement. $\square$

Proposition 5.7. *Let X be a complexity-one T-variety equivariantly covered by toric charts and $\overline{X}$ be a total coordinate space of the Cox ring of X . Then $\overline{X}$ is flexible.*

Proof. Let $\mathbf{k}[\overline{X}]$ be as in Proposition 5.6. We assume that $\overline{X}$ is naturally embedded into an affine space $\mathbb{A}^N = \text{Spec } \mathbf{k}[S_1, \dots, S_N; T_{\ell,j} \mid 0 \leq \ell \leq m, 1 \leq j \leq n_\ell]$.

The images of monomials $A_0, \dots, A_m$ in $\mathbf{k}[\overline{X}]$ span a two-dimensional subspace, and no two of them are collinear. Therefore, we may permute $A_0, \dots, A_m$ along with a proper change of their coefficients, indices of variables, and numbers z_i .

Lemma 5.8. *The point $x \in \overline{X}$ is singular if and only if there are at least three monomials A_i such that all their partial derivatives are vanishing at x .*

Proof. Let x be singular and denote $L_\ell = z_\ell \cdot A_0 + A_1 + A_\ell$ for $\ell = 2, \dots, m$, then there is a non-trivial linear combination $L \in \langle L_2, \dots, L_m \rangle_{\mathbf{k}}$, whose partial derivatives vanish at x . Since L is a sum of at least three monomials A_i , whose partial derivatives also vanish, the statement follows.

Conversely, given three monomials with partial derivatives vanishing at x , we assume that they are A_0, A_1, A_2 . Then we take $L = L_2$. $\square$

So, for a smooth point $x \in X$, up to permutation of monomials and variables we assume that each monomial A_i , $i = 2, \dots, m$, has a non-zero partial derivative, say, $\frac{\partial A_i}{\partial T_{i,1}}(x) \neq 0$. Moreover, we may choose $T_{i,1}$ to be linear in A_i . Indeed, if $T_{i,1}$

is not linear, then $A_i(x) \neq 0$, and we take $T_{i,1}$ to be a linear variable by 5.6.(iv). We denote $B_i = \frac{\partial A_i}{\partial T_{i,1}}$, which is non-zero at x .

Given a set of arbitrary numbers $c_{0,1}, \dots, c_{0,n_0}, c_{1,1}, \dots, c_{1,n_1} \in \mathbf{k}$, we construct a $\mathbb{G}_a$ -action ϕ on $\mathbb{A}^N$ in two steps. First, denoting a parameter of $\mathbb{G}_a$ by t , we let

$$\begin{aligned} \phi^*: \mathbf{k}[\mathbb{A}^N] &\rightarrow \mathbf{k}[\mathbb{A}^N] \otimes \mathbf{k}[t], \\ S_i &\mapsto S_i, & \text{for } i = 1, \dots, n_S, \\ T_{0,j} &\mapsto T_{0,j} + tc_{0,j} \prod_{k=2}^m B_k, & \text{for } j = 1, \dots, n_0, \\ T_{1,j} &\mapsto T_{1,j} + tc_{1,j} \prod_{k=2}^m B_k, & \text{for } j = 1, \dots, n_1. \end{aligned}$$

Then, for some $H_0, H_1 \in \mathbf{k}[\mathbb{A}^N] \otimes \mathbf{k}[\mathbb{G}_a]$,

$$\begin{aligned} \phi^*(A_0) &= A_0 + H_0 \prod_{k=2}^m B_k, \\ \phi^*(A_1) &= A_1 + H_1 \prod_{k=2}^m B_k. \end{aligned}$$

Now, for each $\ell = 2, \dots, m$ we let

$$\begin{aligned} T_{\ell,1} &\mapsto T_{\ell,1} - (z_\ell \cdot H_0 + H_1) \prod_{\substack{2 \leq k \leq m \\ k \neq \ell}} B_k, \\ T_{\ell,j} &\mapsto T_{\ell,j}, & \text{for } j = 2, \dots, n_\ell. \end{aligned}$$

Then the trinomial $z_\ell \cdot A_0 + A_1 + A_\ell$ is fixed by ϕ^* , so ϕ preserves $\overline{X} \subset \mathbb{A}^N$. Thus, we have constructed a $\mathbb{G}_a$ -action on the total coordinate space $\overline{X}$, which we also denote by ϕ .

As said before, for a chosen smooth point x we have $B_i(x) \neq 0$. Let us take another smooth point $y \in \overline{X}$ with non-zero coordinates and move x to y by $\mathbb{G}_a$ -actions, denoting images of x by same letter. By translations along coordinates $S_1, \dots, S_{n_S}$ we ‘equalize’ them, i.e., obtain $S_i(x) = S_i(y)$, $i = 1, \dots, n_S$.

Since $c_{0,1}, \dots, c_{0,n_0}, c_{1,1}, \dots, c_{1,n_1} \in \mathbf{k}$ are arbitrary, with ϕ we also equalize coordinates $T_{0,j}$, $j = 1, \dots, n_0$, and $T_{1,j}$, $j = 1, \dots, n_1$, at x and y . Now, let $T_{1,1}$ be a linear variable in A_1 , then for each $\ell = 2, \dots, m$ we construct a $\mathbb{G}_a$ -action ϕ_ℓ by permuting monomials A_1 and A_ℓ and applying the procedure above. Since $\frac{\partial A_1}{\partial T_{1,1}}(x) = \frac{\partial A_1}{\partial T_{1,1}}(y) \neq 0$, with ϕ_ℓ we may equalize coordinates $T_{\ell,2}, \dots, T_{\ell,m}$, but break the equality of coordinate $T_{1,1}$, which we restore with ϕ .

Proceeding this way for each $\ell = 2, \dots, m$, we equalize all coordinates except $T_{2,1}, \dots, T_{m,1}$. But in this case the equation $z_\ell \cdot A_0 + A_1 + A_\ell = 0$ with condition $B_\ell(x), B_\ell(y) \neq 0$ implies $T_{\ell,1}(x) = T_{\ell,1}(y)$ for each ℓ . So, we may send any smooth to any point with non-zero coordinates, hence the action of $\text{SAut } \overline{X}$ is transitive on smooth points of $\overline{X}$. $\square$

Theorem 5.9. *The total coordinate space of a complete smooth T-variety of complexity one is flexible.*

Proof. By [APS14, Thm A.1], every smooth complete rational T-variety of complexity one is covered by affine spaces, so the statement follows from Proposition 5.7. $\square$

5.3. Flexibility of total coordinate spaces vs. flexible coverings. In [APS14] it was proved that for a variety with an open covering by affine spaces one obtains flexibility of the universal torsor. However, it is not clear whether the flexibility property extends to the total coordinate space. This motivates the following even more general question.

Question 5.10. Provided a variety admits a covering by flexible affine subsets, does this imply flexibility of the total coordinate space?

It is also very tempting to try to connect flexibility of the total coordinate space of the Cox ring of X with that of affine cones over X . The following example shows that flexibility of the total coordinate space does not imply flexibility of all affine cones.

Example 5.11 (Del Pezzo surfaces). We have seen in Section 5.1 that all total coordinate spaces of del Pezzo surfaces are flexible. On the other hand, del Pezzo surfaces are covered by affine spaces, which are flexible. Concerning flexibility of affine cones it was shown in [Per13, PW16] that for degree 4 and 5 all affine cones are flexible, but by [KPZ14, CPW16] the anti-canonical cones over del Pezzo surfaces of degree 3, 2, and 1 are not flexible.

One may still ask if flexibility of all affine cones implies flexibility of the total coordinate space or for a more subtle relation, e.g. involving the grading of the Cox ring.

Question 5.12. Is there a relation between flexibility of the total coordinate space of X and the fact that *all* affine cones over X are flexible?

Let us give some illustrating examples for these questions.

Example 5.13 (Toric varieties). The Cox ring of a toric variety is polynomial. Hence, the total coordinate space is flexible. On the other hand, the torus invariant affine charts and also the affine cones of a toric variety are again toric and hence flexible by [AKZ12, Theorem 0.2].

Example 5.14 (Blowups of $\mathbb{P}^n$ in cubic hypersurfaces inside hyperplanes). The blowup constructions from Example 1.7 give varieties for which all affine cones are flexible, as we have seen. On the other hand, the total coordinate space is flexible, as we see in the following.

We can consider the k^* -action on $\mathbb{P}^n$ given by multiplication with the coordinate x_0 . It comes with a natural quotient map to $\mathbb{P}^{n-1}$ being defined outside the isolated fixed point $(1 : 0 : \dots : 0)$. Then the centers of our blowups are fixed points of the action and we obtain induced actions on X and X' with natural quotient maps given by composition of the original quotient map with the blowup. Now, we may use Theorem 1.2 in [HS10] to calculate the Cox rings

$$\mathcal{R}(X) \cong \mathbf{k}[T_0, \dots, T_n, T'_n, S_1] / (T_n T'_n - f(T_0, \dots, T_{n-1}, 0))$$

and

$$\mathcal{R}(X') \cong \mathbf{k}[T_0, \dots, T_n, T'_n, S_1, S_2] / (T_n T'_n - f(T_0, \dots, T_{n-1}, 0)).$$

We see that they are suspensions over an affine space and hence flexible by Theorem 0.2 in [AKZ12].

Example 5.15 (T-varieties of complexity one). The proof of Theorem 5.9 implies that for T-varieties of complexity one the condition of being covered by toric (and hence flexible) charts is enough to obtain flexibility of the total coordinate space. On the other hand, to conclude flexibility of all affine cones we had to impose the stronger (technical) condition of Theorem 4.4.

REFERENCES

- [AFK⁺13] Ivan Arzhantsev, Hubert Flenner, Shulim Kaliman, Frank Kutzschebauch, and Mikhail Zaidenberg. Flexible varieties and automorphism groups. *Duke Mathematical Journal*, 162(4):767–823, 2013.
- [AH06] Klaus Altmann and Jürgen Hausen. Polyhedral divisors and algebraic torus actions. *Math. Ann.*, 334(3):557–607, 2006.
- [AHS08] Klaus Altmann, Jürgen Hausen, and Hendrik Süß. Gluing affine torus actions via divisorial fans. *Transform. Groups*, 13(2):215–242, 2008.
- [AIP⁺11] Klaus Altmann, Nathan Ilten, Lars Petersen, Hendrik Süß, and Robert Vollmert. The Geometry of T-Varieties. In Piotr Pragacz, editor, *Contributions to Algebraic Geometry*, Impanga Lect Notes, 2011.
- [AKZ12] Ivan Arzhantsev, Karine Kuyumzhyan, and Mikhail Zaidenberg. Flag varieties, toric varieties, and suspensions: Three instances of infinite transitivity. *Sbornik: Mathematics*, 203(7):923–949, 2012.
- [APS14] Ivan Arzhantsev, Alexander Perepechko, and Hendrik Süß. Infinite transitivity on universal torsors. *Journal of the London Mathematical Society*, 89(3):762–778, 2014.
- [BH03] Florian Berchtold and Jürgen Hausen. Homogeneous coordinates for algebraic varieties. *J. Algebra*, 266(2):636–670, 2003.
- [BP04] Victor Batyrev and Oleg Popov. The Cox ring of a del Pezzo surface. In *Arithmetic of higher-dimensional algebraic varieties. Proceedings of the workshop on rational and integral points of higher-dimensional varieties, Palo Alto, CA, USA, December 11–20, 2002*, pages 85–103. Boston, MA: Birkhäuser, 2004.
- [Bro13] Morgan Brown. Singularities of Cox rings of Fano varieties. *Journal de Mathématiques Pures et Appliquées*, 99(6):655–667, 2013.
- [CCM⁺14] Ciro Ciliberto, Maria Angelica Cueto, Massimiliano Mella, Kristian Ranestad, and Piotr Zwiernik. Cremona linearizations of some classical varieties. In *Proceedings of the conference "Homage to Corrado Segre"*, 2014. arXiv preprint arXiv:1403.1814.
- [CLS11] David Cox, John Little, and Hal Schenck. *Toric Varieties*. Graduate studies in mathematics. American Mathematical Soc., 2011.
- [CPW16] Ivan Cheltsov, Jihun Park, and Joonyeong Won. Affine cones over smooth cubic surfaces. *J. Eur. Math. Soc.*, 18:1537–1564, 2016.
- [Der07] Ulrich Derenthal. Universal torsors of del Pezzo surfaces and homogeneous spaces. *Adv. Math.*, 213(2):849–864, 2007.
- [GOST15] Yoshinori Gongyo, Shinnosuke Okawa, Akiyoshi Sannai, and Shunsuke Takagi. Characterization of varieties of Fano type via singularities of Cox rings. *Journal of Algebraic Geometry*, 24(1):159–182, 2015.
- [HS10] Jürgen Hausen and Hendrik Süß. The Cox ring of an algebraic variety with torus action. *Advances in Mathematics*, 225(2):977–1012, 2010.
- [IS11] Nathan Ilten and Hendrik Süß. Polarized complexity-1 T -varieties. *Michigan Math. J.*, 60(3):561–578, 11 2011.
- [IV12] Nathan Ilten and Robert Vollmert. Deformations of rational T -varieties. *J. Algebr. Geom.*, 21(3):531–562, 2012.
- [KPZ11] Takashi Kishimoto, Yuri Prokhorov, and Mikhail Zaidenberg. Group actions on affine cones. In *Montreal Centre de Recherches Mathématiques, CRM Proceedings and Lecture Notes*, volume 54, pages 123–163, 2011.
- [KPZ14] Takashi Kishimoto, Yuri Prokhorov, and Mikhail Zaidenberg. Unipotent group actions on del Pezzo cones. *Algebr. Geom.*, 1(1):46–56, 2014.
- [LS13] Alvaro Liendo and Hendrik Süß. Normal singularities with torus actions. *Tohoku Mathematical Journal*, 65:105–130, January 2013.
- [MM81] Shigefumi Mori and Shigeru Mukai. Classification of fano 3-folds with $b_2 \geq 2$. *manuscripta mathematica*, 36(2):147–162, 1981.
- [MM09] Kayo Masuda and Masayoshi Miyanishi. Lifting of the additive group scheme actions. *Tohoku Math. J. (2)*, 61(2):267–286, 2009.
- [MM15] Laurent Manivel and Mateusz Michałek. Secants of minuscule and cominuscule minimal orbits. *Linear Algebra and its Applications*, 481:288–312, 2015.
- [MOZ15] Mateusz Michałek, Luke Oeding, and Piotr Zwiernik. Secant cumulants and toric geometry. *International Mathematics Research Notices*, 2015(12):4019–4063, 2015.
- [Per13] Alexander Perepechko. Flexibility of affine cones over del pezzo surfaces of degree 4 and 5. *Functional Analysis and Its Applications*, 47(4):284–289, 2013.
- [PS11] Lars Petersen and Hendrik Süß. Torus invariant divisors. *Israel Journal of Mathematics*, 182:481–505, 2011.

- [PW16] Jihun Park and Joonyeong Won. Flexible affine cones over del Pezzo surfaces of degree 4. *Eur. J. Math.*, 2(1):304–318, 2016.
- [Süß14] Hendrik Süß. Fano threefolds with 2-torus action – a picture book. *Documenta Mathematica*, 19:905–914, 2014.
- [SZ13] Bernd Sturmfels and Piotr Zwiernik. Binary cumulant varieties. *Annals of Combinatorics*, 17(1):229–250, 2013.
- [TVV09] Damiano Testa, Anthony Várilly-Alvarado, and Mauricio Velasco. Cox rings of degree one del Pezzo surfaces. *Algebra Number Theory*, 3(7):729–761, 2009.
- [Ver09] Peter Vermeire. Singularities of the secant variety. *Journal of Pure and Applied Algebra*, 213(6):1129–1132, 2009.
- [Zak05] Fedor Zak. *Tangents and secants of algebraic varieties*, volume 127. American Mathematical Soc., 2005.
- [Zwi12] Piotr Zwiernik. L-cumulants, L-cumulant embeddings and algebraic statistics. *Journal of Algebraic Statistics*, 3(1), 2012.
- [Zwi15] Piotr Zwiernik. *Semialgebraic Statistics and Latent Tree Models*. CRC Press, 2015.

POLISH ACADEMY OF SCIENCES, MATHEMATICAL INSTITUT, ŚNIADECKICH 8, 00-956 WARSZAWA, POLAND, FREIE UNIVERSITÄT, BERLIN, GERMANY
E-mail address: `mmichalek@impan.pl`

KHARKEVICH INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, 19 BOLSHOY KARETNY PER., 127994 MOSCOW, RUSSIA
E-mail address: `perepeal@gmail.com`

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF MANCHESTER, ALAN TURING BUILDING, OXFORD ROAD MANCHESTER M13 9PL
E-mail address: `hendrik.suess@manchester.ac.uk`